

Thermodynamic equilibrium and its stability for microcanonical systems described by the Sharma-Taneja-Mittal entropy

A.M. Scarfone^{1,2*} and T. Wada^{1,3†}

¹*Dipartimento di Fisica and ²Istituto Nazionale di Fisica della Materia (INFM), Politecnico di Torino, Corso Duca degli Abruzzi 24, 10129 Torino, Italy.*

³*Department of Electrical and Electronic Engineering, Ibaraki University, Hitachi, Ibaraki, 316-8511, Japan*

(Dated: February 2, 2008)

It is generally assumed that the thermodynamic stability of equilibrium state is reflected by the concavity of entropy. We inquire, in the microcanonical picture, on the validity of this statement for systems described by the two-parametric entropy $S_{\kappa,r}$ of Sharma-Taneja-Mittal. We analyze the “composability” rule for two statistically independent systems, A and B, described by the entropy $S_{\kappa,r}$ with the same set of the deformation parameters. It is shown that, in spite of the concavity of the entropy, the “composability” rule modifies the thermodynamic stability conditions of the equilibrium state. Depending on the values assumed by the deformation parameters, when the relation $S_{\kappa,r}(A \cup B) > S_{\kappa,r}(A) + S_{\kappa,r}(B)$ holds (super-additive systems), the concavity conditions does imply the thermodynamics stability. Otherwise, when the relation $S_{\kappa,r}(A \cup B) < S_{\kappa,r}(A) + S_{\kappa,r}(B)$ holds (sub-additive systems), the concavity conditions does not imply the thermodynamical stability of the equilibrium state.

PACS numbers: 02.50.-r, 05.20.-y, 05.90.+m, 65.40.Gr

Keywords: Generalized entropies, Concavity, Thermodynamic equilibrium, Thermodynamic stability.

I. INTRODUCTION

The MaxEnt principle of the thermodynamics, pioneered by Gibbs and Janes [1], implies that, at equilibrium, both $dS = 0$ and $d^2S < 0$. The first of these conditions states that entropy is an extremum, whereas the second condition states that this extremum is a maximum.

It is well-known that from the second condition follow the concavity conditions for the Boltzmann-Gibbs entropy which are equivalent to the thermodynamic stability conditions of its equilibrium distribution [2, 3]. Some interesting physical implications arise from the thermodynamic stability conditions. For instance, the positivity of the heat capacity assures that, for two bodies in thermal contact and with different temperatures, heat flows from the hot body to the cold one.

In the present days it is widely accepted that the Boltzmann-Gibbs distribution represents only a special case among the great diversity of statistical distributions observed in nature. In many cases such distributions show asymptotic long tails with a power-law behavior. Examples include anomalous diffusion and Levy-flight [4, 5], turbulence [6], self-gravitating systems [7], high T_c superconductivity [8], Bose-Einstein condensation [9], kinetics of charge particles [10], biological systems [11] and others.

To deal with such *anomalous* statistical systems, some generalizations of the well-known Boltzmann-Gibbs en-

trophy have been advanced, with the purpose, by a way, to incorporate the newly observed phenomenologies and, on the other hand, to mimics the beautiful mathematical structure of the standard thermostatics theory [12, 13, 14]. A possible way to do this is to replace the standard logarithm in the Boltzmann-Gibbs entropy, $S(p) = -k_B \sum_i p_i \ln(p_i)$, with its generalized version [15, 16].

In this work we investigate the relationship between the concavity conditions and the thermodynamic stability conditions of the equilibrium distribution of a conservative system, with fixed energy and volume, described by a generalized entropy.

A preliminary investigation along this question can be found in Refs. [17, 18, 19, 20].

As a working tool we employ the two-parameter entropy of Sharma-Taneja-Mittal [21, 22, 23, 24]. Although, an entropy containing two free parameters could sound unlike on the physical ground, the Sharma-Taneja-Mittal entropy includes, as special cases, some one-parameter entropies already proposed in literature, like the Tsallis entropy [25, 26], the Abe entropy [27, 28] and the Kaniadakis entropy [29, 30]. Consequently, Sharma-Taneja-Mittal entropy enables us to consider all this one-parameter entropies in a unified scheme.

In Refs. [31, 32] it has been addressed the question on the existence of a generalized trace-form entropy

$$S(p) = - \sum_{i=1}^W p_i \Lambda(p_i), \quad (1.1)$$

(throughout this paper we use units with the Boltzmann constant $k_B = 1$) preserving unaltered the epistemological structure of the standard statistical mechanics. In Eq. (1.1) $\Lambda(x)$ is a deformed logarithm [16] replacing the

*Electronic address: antonio.scarfone@polito.it

†Electronic address: wada@ee.ibaraki.ac.jp

standard one, $\{p_i\}_{i=1,\dots,W}$ is a discrete probability distribution function and W the number of microscopically accessible states. By requiring that the entropy (1.1) preserves the mathematical properties physically motivated [30], the following differential-functional equation has been obtained

$$\frac{d}{dp_j} [p_j \Lambda(p_j)] = \lambda \Lambda\left(\frac{p_j}{\alpha}\right). \quad (1.2)$$

A physically suitable solution $\Lambda(x) \equiv \ln_{\{\kappa, r\}}(x)$ can be written as

$$\ln_{\{\kappa, r\}}(x) = x^r \frac{x^\kappa - x^{-\kappa}}{2\kappa}, \quad (1.3)$$

which satisfies Eq. (1.2) with

$$\alpha = \left(\frac{1+r-\kappa}{1+r+\kappa} \right)^{1/2\kappa}, \quad (1.4)$$

and

$$\lambda = \frac{(1+r-\kappa)^{(r+\kappa)/2\kappa}}{(1+r+\kappa)^{(r-\kappa)/2\kappa}}. \quad (1.5)$$

Taking into account of Eq. (1.3), the generalized entropy (1.1) assumes the form

$$\begin{aligned} S_{\kappa, r}(p) &= - \sum_{i=1}^W (p_i)^{r+1} \frac{(p_i)^\kappa - (p_i)^{-\kappa}}{2\kappa} \\ &= - \sum_{i=1}^W p_i \ln_{\{\kappa, r\}}(p_i), \end{aligned} \quad (1.6)$$

which has been introduced for the first time in Refs. [21, 22, 23] and successively reconsidered in Ref. [24].

Eq. (1.6) mimics the expression of the Boltzmann-Gibbs entropy by replacing the standard logarithm $\ln(x)$ with the two-parametric deformed logarithm $\ln_{\{\kappa, r\}}(x)$.

The distribution obtained by optimizing Eq. (1.6), under standard linear energy constraint $\sum_i p_i E_i = \langle E \rangle$ and the normalization constraint $\sum_i p_i = 1$, assumes the form

$$p_i = \alpha \exp_{\{\kappa, r\}} \left(-\frac{\beta}{\lambda} (E_i - \mu) \right), \quad (1.7)$$

where $\exp_{\{\kappa, r\}}(x)$ is the inverse function of $\ln_{\{\kappa, r\}}(x)$, namely the deformed exponential.

Remarkably, Eq. (1.7) exhibits an asymptotic power law behavior, with $p_i \sim E_i^{1/(r-\kappa)}$, for large E_i . This entropy possesses positivity, continuity, symmetry, expandibility, decisivity, maximality, concavity, and is Lesche stable whenever $(\kappa, r) \in \mathcal{R}$, where the two dimensional region $\mathcal{R}$ is defined by $-|\kappa| \leq r \leq |\kappa|$ for $|\kappa| < 1/2$ and $|\kappa| - 1 < r < 1 - |\kappa|$ for $1/2 \leq |\kappa| < 1$.

We remark that the deformed logarithm (1.3) reduces to the standard logarithm in the $(\kappa, r) \rightarrow (0, 0)$ limit ($\ln_{\{0, 0\}}(x) \equiv \ln x$) and, in the same limit, Eq. (1.6) reduces to the Boltzmann-Gibbs entropy. [Refer to the

Appendix A for the main mathematical properties of the deformed logarithm (1.3)].

The plan of the paper is the following. In the next Section 2, we consider the Sharma-Taneja-Mittal entropy and its distribution in the microcanonical framework. In Section 3 we derive the ‘‘composability’’ rule for two statistically independent systems A and B with the same set of the deformation parameters. In Section 4 we inquire on the functional relationship between the Sharma-Taneja-Mittal entropy and the definitions of temperature and pressure obtained as equivalence relations at the equilibrium configuration. In Section 5 we examine the thermodynamic response produced by perturbing the system away from the equilibrium. The perturbations are generated by repartitioning the energy (heat transfer) and the volume (work transfer) between the two systems A and B. According to the MaxEnt principle such processes lead to a lower entropy, provided that the whole system AUB was initially in a stable equilibrium. By analyzing the signs of the entropy changes for these processes we obtain the corresponding thermodynamic stability conditions. Finally, in section 6 we relate this results to some known one-parameter cases. Concluding remarks are reported in section 7. In Appendix A we give some mathematical properties of the deformed logarithm, and Appendix B deals with the sketch of some proofs.

II. MICROCANONICAL SHARMA-TANEJA-MITTL ENTROPY

According to the MaxEnt principle, the equilibrium distribution is the one that maximizes the entropy under the constraints imposed on the probability distribution. In the microcanonical picture, the system has fixed total energy E and volume V , and the distribution $p \equiv \{p_i\}_{i=1,\dots,W}$ is obtained by optimizing the entropy (1.6) under the only constraint on the normalization

$$\sum_{i=1}^W p_i = 1. \quad (2.1)$$

Thus, we have to deal with the variational problem

$$\frac{\delta}{\delta p_j} \left(S_{\kappa, r}(p) - \gamma \sum_{i=1}^W p_i \right) = 0, \quad (2.2)$$

where γ is the Lagrange multiplier associated with the constraint (2.1). By taking into account Eqs. (1.1)-(1.2), it follows

$$\lambda \ln_{\{\kappa, r\}} \left(\frac{p_j}{\alpha} \right) + \gamma = 0, \quad (2.3)$$

and by means of the deformed exponential $\exp_{\{\kappa, r\}}(x)$, we obtain

$$p_j = \alpha \exp_{\{\kappa, r\}} \left(-\frac{\gamma}{\lambda} \right). \quad (2.4)$$

Since this distribution does not depend on the index j , according to Eq. (2.1), it takes the form

$$p_j = \frac{1}{W(E, V)}, \quad \text{with} \quad j = 1, \dots, W, \quad (2.5)$$

where we took into account that the number of accessible states $W(E, V)$ is a function of the energy E and the volume V of the system.

By substituting Eq. (2.5) into Eq. (1.6) we obtain its expression in the microcanonical picture

$$\begin{aligned} S_{\kappa, r}(E, V) &= -\ln_{\{\kappa, r\}} \left(\frac{1}{W(E, V)} \right) \\ &= \ln_{\{\kappa, -r\}} (W(E, V)). \end{aligned} \quad (2.6)$$

This is evocative of the well-known Boltzmann formula $S = \ln(W)$, which is indeed recovered in the $(\kappa, r) \rightarrow (0, 0)$ limit.

We observe that the concavity of the function $\ln_{\{\kappa, r\}}(x)$ w.r.t. its argument x does not necessarily imply the concavity of the entropy (2.6) w.r.t E and V . The concavity conditions for the given problem follow from the analysis of the sign of the eigenvalues of the Hessian matrix associated to Eq. (2.6). In particular, by requiring that the following quadratic form is negative definite [33]

$$\phi(\mathbf{y}; E, V) = \frac{\partial^2 S_{\kappa, r}}{\partial E^2} y_E^2 + 2 \frac{\partial^2 S_{\kappa, r}}{\partial E \partial V} y_E y_V + \frac{\partial^2 S_{\kappa, r}}{\partial V^2} y_V^2, \quad (2.7)$$

for any arbitrary vector $\mathbf{y} \equiv (y_E, y_V)$, we obtain the following relations

$$\frac{\partial^2 S_{\kappa, r}}{\partial E^2} < 0, \quad (2.8)$$

and

$$\frac{\partial^2 S_{\kappa, r}}{\partial E^2} \frac{\partial^2 S_{\kappa, r}}{\partial V^2} - \left(\frac{\partial^2 S_{\kappa, r}}{\partial E \partial V} \right)^2 > 0, \quad (2.9)$$

stating the concavity conditions for the entropy (2.6).

III. COMPOSED SYSTEMS

Let us consider two systems A and B described by the entropy (2.6), with the same set of the deformation parameters.

We denote with $W_A \equiv W(E_A, V_A)$ and $W_B \equiv W(E_B, V_B)$ the number of accessible states of the two systems A and B respectively and hypothesize a statistical independence of A and B, in the sense that the number of accessible states $W_{A \cup B} \equiv W(E_{A \cup B}, V_{A \cup B})$ of the composed system AUB is given by $W_{A \cup B} = W_A W_B$. In Ref. [34] the most general form of pseudoadditivity of

composable entropies, as prescribed by the existence of equilibrium, has been obtained. The main result reads

$$H[S(A \cup B)] = H[S(A)] + H[S(B)] + \lambda H[S(A)] H[S(B)], \quad (3.1)$$

where $H(x)$ is a certain differentiable function, λ denotes the set of deformation parameters, whilst $S(A)$, $S(B)$ and $S(A \cup B)$ are the entropies of the systems A, B and AUB, respectively.

It is easy to show that Eq. (3.1) is fulfilled by the entropy (2.6) if we define

$$\begin{aligned} H[S_{\kappa, r}(W)] &= -S_{\kappa, r}(W) \left[\exp_{\{\kappa, r\}} (-S_{\kappa, r}(W)) \right]^{-r-\kappa} \\ &= W^{r+\kappa} \ln_{\{\kappa, r\}} \left(\frac{1}{W} \right). \end{aligned} \quad (3.2)$$

In fact, by using Eq. (A.10) given in Appendix A, we have

$$\begin{aligned} &(W_A W_B)^{r+\kappa} \ln_{\{\kappa, r\}} \left(\frac{1}{W_A W_B} \right) \\ &= (W_A)^{r+\kappa} \ln_{\{\kappa, r\}} \left(\frac{1}{W_A} \right) + (W_B)^{r+\kappa} \ln_{\{\kappa, r\}} \left(\frac{1}{W_B} \right) \\ &\quad - 2\kappa \left[(W_A)^{r+\kappa} \ln_{\{\kappa, r\}} \left(\frac{1}{W_A} \right) \right] \\ &\quad \times \left[(W_B)^{r+\kappa} \ln_{\{\kappa, r\}} \left(\frac{1}{W_B} \right) \right], \end{aligned} \quad (3.3)$$

which has the same structure of Eq. (3.1) with $H[S_{\kappa, r}]$ given in Eq. (3.2) and $\lambda = -2\kappa$.

After multiplying Eq. (3.3) by $(W_A W_B)^{-r-\kappa}$ and recalling the invariance of the entropy (2.6) under the interchange of $\kappa \leftrightarrow -\kappa$, Eq. (3.3) becomes

$$S_{\kappa, r}(A \cup B) = S_{\kappa, r}(A) \mathcal{I}_{\kappa, r}(B) + \mathcal{I}_{\kappa, r}(A) S_{\kappa, r}(B), \quad (3.4)$$

where the function $\mathcal{I}_{\kappa, r}(p)$, for a given distribution function $p = \{p_i\}_{i=1, \dots, W}$, is defined by

$$\mathcal{I}_{\kappa, r}(p) = \sum_{i=1}^W (p_i)^{r+1} \frac{(p_i)^\kappa + (p_i)^{-\kappa}}{2}, \quad (3.5)$$

with $\mathcal{I}_{0,0}(p) = 1$ and reduces, for the uniform distribution (2.5), to

$$\mathcal{I}_{\kappa, r} \left(\frac{1}{W(E, V)} \right) = \frac{[W(E, V)]^{-r-\kappa} + [W(E, V)]^{-r+\kappa}}{2}. \quad (3.6)$$

We remark that Eq. (3.6) actually is a function of the entropy (2.6) through the relation

$$\mathcal{I}_{\kappa, r}(x) = \kappa S_{\kappa, r}(x) + \left[\exp_{\{\kappa, r\}} (-S_{\kappa, r}(x)) \right]^{r+\kappa}. \quad (3.7)$$

It is worthy to observe that Eq. (3.4) still holds, for a canonical distribution, also if the entropy (1.6), in this case, does not satisfy the criteria dictated by Eq. (3.1).

Eq. (3.4) expresses the “composability” properties for a system described by the entropy (2.6), and, in the $(\kappa, r) \rightarrow (0, 0)$ limit, we recover the well-known additivity rule of the Boltzmann entropy $S(A \cup B) = S(A) + S(B)$.

In the following we analyze in more details this equation. According to the results given in the Appendix A, when $(\kappa, r) \in \mathcal{R}|_{r \leq 0}$ it follows $\mathcal{I}_{\kappa, r}(1/W) > 1$ for $W > 1$. Consequently, from Eq. (3.4) we obtain

$$S_{\kappa, r}(A \cup B) > S_{\kappa, r}(A) + S_{\kappa, r}(B), \quad (3.8)$$

and the entropy (2.6) exhibits a super-additive behavior. The analysis of the Eq. (3.4) becomes more complicated in the complementary region $\mathcal{R}|_{r > 0}$. In fact, for $r > 0$ there exists a threshold point $W_t(\kappa, r) > 1$, which is defined by

$$\mathcal{I}_{\kappa, r}\left(\frac{1}{W_t}\right) = 1, \quad (3.9)$$

so that $\mathcal{I}_{\kappa, r}(1/W) \leq 1$ when $1 < W \leq W_t$ whereas $\mathcal{I}_{\kappa, r}(1/W) > 1$ when $W > W_t$. Consequently, for $(\kappa, r) \in \mathcal{R}|_{r > 0}$ we have a sub-additive behavior

$$S_{\kappa, r}(A \cup B) < S_{\kappa, r}(A) + S_{\kappa, r}(B), \quad (3.10)$$

when both $1 < W_A < W_t$ and $1 < W_B < W_t$, whereas the super-additive behavior (3.8) is recovered whenever $W_A > W_t$ and $W_B > W_t$. In the intermediate situation $1 < W_A < W_t$ and $W_B > W_t$, or $W_A > W_t$ and $1 < W_B < W_t$, the character of the composition law is not well determined, depending on the values of W_A and W_B . Thus, for $(\kappa, r) \in \mathcal{R}|_{r > 0}$ the value of the entropy of a composed system $A \cup B$, w.r.t. the sum of the entropies of the two separate systems A and B, depends on the *size* of the two systems [55]. *Small* systems exhibit a sub-additive behavior, which becomes super-additive when both the systems grow over the threshold point W_t .

As consequence, super-additivity behavior emerges with *larger* systems.

We observe that the threshold point W_t becomes larger and larger, for $r \rightarrow \kappa$, according to

$$\lim_{r \rightarrow \kappa} W_t(\kappa, r) \rightarrow \infty. \quad (3.11)$$

As a consequence, $\mathcal{I}_{\kappa, \kappa} \leq 1$ and the entropy $S_{\kappa, \kappa}(A \cup B)$ has always a sub-additive behavior.

IV. THERMAL AND MECHANICAL EQUILIBRIUM

Possible definitions of temperature and pressure, in the construction of a generalized framework of thermodynamics, have been proposed in Refs. [35, 36, 37] through the study of the equilibrium configuration. Such method can be successfully applied to the generalized entropy under inspection.

We assume that both energy and volume are additive quantities, i.e., $E_{A \cup B} = E_A + E_B$ and $V_{A \cup B} = V_A + V_B$. A different approach, by utilizing nonadditive energy and volume, within the framework of nonextensive statistical mechanics, has been explored in Ref. [38].

Let us consider an isolated system $A \cup B$ composed by two, statistically independent, systems A and B in contact through an ideal wall. The wall permits transfer of energy (heat) and/or volume (work) between the two systems but is adiabatic with respect to any other interaction.

We suppose that the system, initially at the thermal and mechanical equilibrium, undergoes a small fluctuation of energy and volume between A and B. According to the MaxEnt principle the variation of the entropy evaluated at the first order in δE and δV must vanish

$$\delta S_{\kappa, r}(A \cup B) = 0, \quad (4.1)$$

where

$$\delta(E_A + E_B) = 0, \quad (4.2)$$

$$\delta(V_A + V_B) = 0. \quad (4.3)$$

From Eq. (3.4) we obtain (see Appendix B)

$$\frac{1}{\mathcal{I}_{\kappa, r} - r S_{\kappa, r}} \frac{\partial S_{\kappa, r}}{\partial E} \Big|_A = \frac{1}{\mathcal{I}_{\kappa, r} - r S_{\kappa, r}} \frac{\partial S_{\kappa, r}}{\partial E} \Big|_B, \quad (4.4)$$

and

$$\frac{1}{\mathcal{I}_{\kappa, r} - r S_{\kappa, r}} \frac{\partial S_{\kappa, r}}{\partial V} \Big|_A = \frac{1}{\mathcal{I}_{\kappa, r} - r S_{\kappa, r}} \frac{\partial S_{\kappa, r}}{\partial V} \Big|_B. \quad (4.5)$$

Actually Eqs. (4.4) and (4.5) state the analytical formulation of the zeroth law of the thermodynamics for the system under inspection and define, as equivalence relations, modulo of a multiplicative constant, the temperature

$$\frac{1}{T} = \frac{1}{\mathcal{I}_{\kappa, r} - r S_{\kappa, r}} \frac{\partial S_{\kappa, r}}{\partial E}, \quad (4.6)$$

and the pressure

$$P = \frac{T}{\mathcal{I}_{\kappa, r} - r S_{\kappa, r}} \frac{\partial S_{\kappa, r}}{\partial V}, \quad (4.7)$$

which, accounting Eq. (3.7), are given only through the entropy $S_{\kappa, r}$.

The standard relations of the classical thermodynamics

$$\frac{1}{T} = \frac{\partial S}{\partial E}, \quad (4.8)$$

and

$$P = T \frac{\partial S}{\partial V}, \quad (4.9)$$

are recovered in the $(\kappa, r) \rightarrow (0, 0)$ limit.

It is worth to observe that, by using Eq. (2.6), Eqs. (4.6)-(4.7) can be written as

$$\frac{1}{T} = \frac{\partial}{\partial E} \ln(W), \quad (4.10)$$

and

$$P = T \frac{\partial}{\partial V} \ln(W), \quad (4.11)$$

which are positive definite quantities if $W(E, V)$ is a monotonic increasing function with respect to both E and V .

In Refs. [39, 40] has been noted that in the micro-canonical framework of the Tsallis' thermostatics, the definitions of T and P , obtained through the study of the equilibrium configuration, lead to expressions which coincide with those obtained by using the standard Boltzmann formalism of statistical mechanics. This results still hold in presence of the entropy (2.6), as can be seen from Eqs. (4.10) and (4.11), which define temperature and pressure in function of W and coincide with the standard definitions adopted in the Boltzmann theory.

V. THERMODYNAMIC STABILITY

In this section we examine the thermodynamic response produced by perturbing the system which is assumed initially in equilibrium. By analyzing the signs of thermodynamic changes, we obtain the thermodynamic stability conditions.

Let us consider a small perturbation of the system through a transfer of an amount of energy and/or volume between A and B: $S_{\kappa, r}(A \cup B) \rightarrow S_{\kappa, r}((A + \delta A) \cup (B + \delta B))$.

According to the MaxEnt principle, such a perturbation leads the system in a new state with a lower entropy

$$S_{\kappa, r}(A \cup B) > S_{\kappa, r}((A + \delta A) \cup (B + \delta B)). \quad (5.1)$$

In Eq. (5.1) we denote $S_{\kappa, r}(A \cup B) \equiv S_{\kappa, r}(E_A + E_B, V_A + V_B)$ and $S_{\kappa, r}((A + \delta A) \cup (B + \delta B)) \equiv S_{\kappa, r}(E_A + \delta E_A, V_A + \delta V_A) \cup (E_B + \delta E_B, V_B + \delta V_B)$ where $\delta E_A = -\delta E_B \equiv \delta E$ and $\delta V_A = -\delta V_B \equiv \delta V$.

Recalling Eq. (3.4), Eq. (5.1) can be written in

$$\begin{aligned} & S_{\kappa, r}(A) \mathcal{I}_{\kappa, r}(B) + \mathcal{I}_{\kappa, r}(A) S_{\kappa, r}(B) > \\ & S_{\kappa, r}(A + \delta A) \mathcal{I}_{\kappa, r}(B + \delta B) \\ & + \mathcal{I}_{\kappa, r}(A + \delta A) S_{\kappa, r}(B + \delta B), \end{aligned} \quad (5.2)$$

and after expanding the r.h.s of Eq. (5.2) up to the second order in δE and δV , we obtain (see the Appendix B)

$$\begin{aligned} & \frac{1}{2} (\mathcal{I}_{\kappa, r} - r S_{\kappa, r}) \Big|_{A \cup B} \\ & \left[\frac{S_{EE} (\delta E)^2 + 2 S_{EV} \delta E \delta V + S_{VV} (\delta V)^2}{\mathcal{I}_{\kappa, r} - r S_{\kappa, r}} \right] \Big|_A \end{aligned}$$

$$+ \frac{S_{EE} (\delta E)^2 + 2 S_{EV} \delta E \delta V + S_{VV} (\delta V)^2}{\mathcal{I}_{\kappa, r} - r S_{\kappa, r}} \Big|_B < 0, \quad (5.3)$$

where we have posed

$$\mathcal{S}_{XY} = \frac{\partial^2 S_{\kappa, r}}{\partial X \partial Y} - \frac{(\kappa^2 + r^2) S_{\kappa, r} - 2 r \mathcal{I}_{\kappa, r}}{(\mathcal{I}_{\kappa, r} - r S_{\kappa, r})^2} \frac{\partial S_{\kappa, r}}{\partial X} \frac{\partial S_{\kappa, r}}{\partial Y}. \quad (5.4)$$

Eq. (5.3) is fulfilled if the following inequalities

$$\mathcal{S}_{EE} < 0, \quad (5.5)$$

$$\mathcal{S}_{EE} \mathcal{S}_{VV} - \mathcal{S}_{EV}^2 > 0, \quad (5.6)$$

are separately satisfied by both systems A and B.

Remark that Eqs. (5.5)-(5.6) have the same structure of Eq. (2.8)-(2.9).

Explicitly, Eqs. (5.5)-(5.6) read

$$\frac{\partial^2 S_{\kappa, r}}{\partial E^2} < \mathcal{A}_{\kappa, r} \left(\frac{\partial S_{\kappa, r}}{\partial E} \right)^2, \quad (5.7)$$

and

$$\frac{\partial^2 S_{\kappa, r}}{\partial E^2} \frac{\partial^2 S_{\kappa, r}}{\partial V^2} - \left(\frac{\partial^2 S_{\kappa, r}}{\partial E \partial V} \right)^2 > \mathcal{A}_{\kappa, r} \mathcal{B}_{\kappa, r} \quad (5.8)$$

where

$$\mathcal{A}_{\kappa, r} = \frac{(\kappa^2 + r^2) S_{\kappa, r} - 2 r \mathcal{I}_{\kappa, r}}{(\mathcal{I}_{\kappa, r} - r S_{\kappa, r})^2}, \quad (5.9)$$

and

$$\begin{aligned} \mathcal{B}_{\kappa, r} = & \left(\frac{\partial^2 S_{\kappa, r}}{\partial E^2} \right)^{-1} \left\{ \left(\frac{\partial^2 S_{\kappa, r}}{\partial E^2} \frac{\partial S_{\kappa, r}}{\partial V} - \frac{\partial^2 S_{\kappa, r}}{\partial E \partial V} \frac{\partial S_{\kappa, r}}{\partial E} \right)^2 \right. \\ & \left. + \left(\frac{\partial S_{\kappa, r}}{\partial E} \right)^2 \left[\frac{\partial^2 S_{\kappa, r}}{\partial E^2} \frac{\partial^2 S_{\kappa, r}}{\partial V^2} - \left(\frac{\partial^2 S_{\kappa, r}}{\partial E \partial V} \right)^2 \right] \right\}. \end{aligned} \quad (5.10)$$

In particular, the quantity $\mathcal{B}_{\kappa, r}$ is negative definite for a concave entropy, as a consequence of Eqs. (2.8)-(2.9).

From Eqs. (5.5) and (5.6) it is trivial to obtain the further inequality $\mathcal{S}_{VV} < 0$, which can be written in

$$\frac{\partial^2 S_{\kappa, r}}{\partial V^2} < \mathcal{A}_{\kappa, r} \left(\frac{\partial S_{\kappa, r}}{\partial V} \right)^2. \quad (5.11)$$

Eqs. (5.7), (5.8) and (5.11) are the announced thermodynamic stability conditions for the family of entropies given in Eq. (1.6), and reduce to Eq. (2.8)-(2.9) in the $(\kappa, r) \rightarrow (0, 0)$ limit. Depending on the sign of the function $\mathcal{A}_{\kappa, r}$ this inequalities are satisfy if the concavity conditions does.

We observe that the equation $\mathcal{A}_{\kappa, r}(\tilde{x}_t) = 0$ has the unique solution given by

$$\tilde{x}_t = \left(\frac{\kappa - r}{\kappa + r} \right)^{1/\kappa}. \quad (5.12)$$

By inspection it follows that $\mathcal{A}_{\kappa,r}(x) > 0$ when $0 \leq x \leq \tilde{x}_t$ whilst $\mathcal{A}_{\kappa,r}(x) < 0$ when $\tilde{x}_t \leq x \leq +\infty$. In the parametric region $(\kappa, r) \in \mathcal{R}|_{r \leq 0}$, accounting for Eq. (5.12), it follows that $1/\tilde{W}_t = \tilde{x}_t > 1$. Thus, being $W \geq 1$, it follows $1/W < 1 < 1/\tilde{W}_t$ so that $\mathcal{A}_{\kappa,r} > 0$ and consequently Eqs. (5.7), (5.8) and (5.11) are fulfilled if the concavity conditions are accomplished.

Differently, in the parametric region $(\kappa, r) \in \mathcal{R}_{r>0}$, we have $1/\tilde{W}_t < 1$. In this case $\mathcal{A}_{\kappa,r} \geq 0$ when $W \geq \tilde{W}_t$. As a consequence we obtain that the concavity conditions imply the thermodynamic stability conditions if and only if both $W_A \geq \tilde{W}_t$ and $W_B \geq \tilde{W}_t$ are satisfied.

At this point we observe that, when $r > 0$

$$\mathcal{I}_{\kappa,r} \left(\frac{1}{\tilde{W}_t} \right) < 1, \quad (5.13)$$

so that

$$\tilde{W}_t < W_t, \quad (5.14)$$

where W_t is the threshold point defined in Eq. (3.9), and the system becomes super-additive when the number of accessible states W_A and W_B beyond W_t . Thus, we can conclude that, whenever the system exhibits a super-additive behavior, the concavity conditions are sufficient to guarantee the thermodynamic stability of the equilibrium configuration.

VI. EXAMPLES

In this section we specify our results to some one-parameter entropies, already known in literature, and belonging to the family of the Sharma-Taneja-Mittal entropy.

In figure 1 we depict the log-linear plots for the three one-parameter entropies discussed in this section, for different values of the deformation parameter. The solid line shows the Boltzmann entropy.

A. Tsallis entropy

As a first example, we consider the Tsallis entropy [25]

$$S_{2-q} = - \sum_{i=1}^W \frac{p_i^{2-q} - p_i}{q-1} = - \sum_{i=1}^W p_i \ln_q(p_i), \quad (6.1)$$

with $0 < q < 2$, which follows from Eq. (1.6) by posing $r = \pm|\kappa|$ and introducing the parameter $q = 1 \mp 2|\kappa|$. We remark that Eq. (6.1) differs from the usual definition adopted in the Tsallis framework which is recovered by replacing $q \rightarrow 2-q$.

In Eq. (6.1) the q -logarithm, $\ln_q(x)$, is defined by

$$\ln_q(x) = \frac{x^{1-q} - 1}{1-q}, \quad (6.2)$$




FIG. 1: Log-linear plot of some one-parameter entropies (in arbitrary units) belonging to the family (2.6) for several values of the deformation parameter: **(A)** Tsallis entropy, [25]; **(B)** Abe entropy, [27]; and **(C)** Kaniadakis entropy, [30]. The solid curves show the Boltzmann entropy.

whereas, its inverse function, namely the q -exponential, is given by

$$\exp_q(x) = [1 + (1-q)x]^{1/(1-q)}. \quad (6.3)$$

Both Eqs. (6.2) and (6.3) fulfill the relations

$$\exp_q(x) \exp_q(y) = \exp_q(x \oplus_q y), \quad (6.4)$$

$$\ln_q(xy) = \ln_q(x) \oplus_q \ln_q(y), \quad (6.5)$$

where the q -deformed sum, introduced in [41, 42], is defined as

$$x \oplus_q y = x + y + (1 - q)xy. \quad (6.6)$$

Eqs. (6.4) and (6.5) reduce, in the $q \rightarrow 1$ limit, to the well known standard relations $\exp(x) \exp(y) = \exp(x+y)$ and $\ln(xy) = \ln(x) + \ln(y)$, respectively, according to $x \oplus_1 y = x + y$.

After its introduction in 1988, Tsallis entropy has been largely applied, as a paradigm, in the study of complex systems having a probability distribution function with a power law behavior in the tail. Typically, these systems are characterized by long-range interactions or long time memory effects that establish a space-time interconnection by the parts which causes a strong interdependence and the existence of a rich structure over several scales. All these induce correlations between the parts of the system which gives origin to a dynamical equilibrium rather than a static equilibrium: the system remains in a metastable configuration that could persist for a long period of time as compared with the characteristic time scale of the underlying microscopic dynamical process.

As it is known, the Tsallis entropy exhibits many interesting proprieties which make it a suitable substitute of the Boltzmann Gibbs entropy in the study of these anomalous systems. Among them we recall that it is concave for $q > 0$, Lesche stable [43], a basic propriety which must be satisfied in order to represent a well defined physical observable [44], and fulfill the Pesin identity [45] stating a relation between the sensitivity to the initial conditions and the (finite) entropy production per unit time. Many other physical proprieties about the Tsallis entropy can be found in [46].

In the microcanonical picture, with a uniform distribution $p_i = 1/W$, Eq. (6.1) reduces to

$$S_{2-q} = -\ln_q \left(\frac{1}{W} \right), \quad (6.7)$$

and we introduce the function $\mathcal{I}_{2-q}$ which, according to Eq. (3.7), can be expressed as a linear function of the entropy

$$\mathcal{I}_{2-q} = \frac{1}{2}(q-1)S_{2-q} + 1. \quad (6.8)$$

From Eq. (6.8) it readily follows that $\mathcal{I}_{2-q} > 1$ for $q > 1$ and $\mathcal{I}_{2-q} < 1$ for $q < 1$, and from Eq. (3.4) we obtain the well-known “composability” rule

$$S_{2-q}(A \cup B) = S_{2-q}(A) + S_{2-q}(B) + (q-1)S_{2-q}(A)S_{2-q}(B), \quad (6.9)$$

which shows that S_{2-q} is sub-additive for $q \in (0, 1)$ and super-additive for $q \in (1, 2)$. Remark that Eq. (6.9) can be readily obtained also from the proprieties (6.5) of the q -logarithm.

Temperature and pressure are defined through [36, 37]

$$\frac{1}{T} = \frac{1}{1 + (q-1)S_{2-q}} \frac{\partial S_{2-q}}{\partial E}, \quad (6.10)$$

$$P = \frac{T}{1 + (q-1)S_{2-q}} \frac{\partial S_{2-q}}{\partial V}, \quad (6.11)$$

respectively, whereas the thermodynamic stability conditions are obtained through Eqs. (5.7)-(5.11) and read [19, 20]

$$\frac{\partial^2 S_{2-q}}{\partial E^2} < \mathcal{A}_{2-q} \left(\frac{\partial S_{2-q}}{\partial E} \right)^2, \quad (6.12)$$

$$\frac{\partial^2 S_{2-q}}{\partial E^2} \frac{\partial^2 S_{2-q}}{\partial V^2} - \left(\frac{\partial^2 S_{2-q}}{\partial E \partial V} \right)^2 > \mathcal{A}_{2-q} \mathcal{B}_{2-q} \quad (6.13)$$

where

$$\mathcal{A}_{2-q} = \frac{q-1}{1 + (q-1)S_{2-q}}. \quad (6.14)$$

Eq. (6.14) is a positive quantity for $q \in (1, 2)$. Consequently, it follows that both Eqs. (6.12) and (6.13) are fulfilled if the concavity conditions are satisfied. Differently, for $q \in (0, 1)$ Eq. (6.14) is a negative quantity. For this range of values of q the thermodynamical stability of the equilibrium configuration does not follow merely from the concavity conditions of S_{2-q} .

Such conclusion is in accordance with the results discussed in Ref. [47].

In spite of the success obtained by the Tsallis entropy in the study of anomalous systems, others entropic forms, with probability distribution function exhibiting an asymptotic power law behavior, have been proposed by different authors. Some of them belong to the family of the Sharma-Taneja-Mittal entropy and we explore them in the next examples.

B. Abe entropy

In Ref. [27] it has been presented a new entropy containing the quantum group deformation structure, through the requirement of the invariance under the interchange $q \leftrightarrow q^{-1}$. This can be accomplished by posing $r = \sqrt{1 + \kappa^2} - 1 > 0$ and $q_A = \sqrt{1 + \kappa^2} + |\kappa|$, so that Eq. (1.6) becomes

$$S_{q_A} = - \sum_{i=1}^W \frac{p_i^{q_A^{-1}} - p_i^{q_A}}{q_A^{-1} - q_A} = - \sum_{i=1}^W p_i \ln_{q_A} (p_i), \quad (6.15)$$

with $1/2 < q_A \leq 2$, and

$$\ln_{q_A}(x) = \frac{x^{(q_A^{-1})-1} - x^{q_A-1}}{q_A^{-1} - q_A}. \quad (6.16)$$

We remark that the inverse function of Eq. (6.16), namely $\exp_{q_A}(x)$, exists because Eq. (6.16) is a monotonic function, but its expression cannot be given in term of elementary functions.

Entropy (6.15) has been applied in [28] to the generalized statistical mechanics study of q -deformed oscillators.

The basic idea is to incorporate the nonadditive feature of the energies of the systems having the quantum group structures with generalized statistics. It has been shown that for large value of $\partial S_{q_A}/\partial E$ the deformation of the entropy gives rise to significative deviations of the Planck distribution with respect to the standard (undeformed) behavior.

In Ref. [27] it has been shown that Abe' entropy can be expressed as a combination of Tsallis' entropy with different deformation parameters. Consequently, many physical proprieties of the former follows from the physical proprieties of the latter. In particular, it can be shown that it is Lesche stable [48], and fulfills the Pesin equality [49].

In the microcanonical picture, the entropy (6.15) becomes

$$S_{q_A} = -\ln_{q_A} \left(\frac{1}{W} \right), \quad (6.17)$$

and we introduce the function $\mathcal{I}_{q_A}$, through Eq. (3.6), which assumes the expression

$$\mathcal{I}_{q_A} = \frac{1}{2} \left(W^{1-(q_A^{-1})} + W^{1-q_A} \right). \quad (6.18)$$

We recall that, according with Eq. (3.7), Eq. (6.18) is a function of the entropy S_{q_A} .

By taking into account the results of appendix A, we have $1/2 \leq \mathcal{I}_{q_A} \leq +\infty$, depending on the value of W . After introducing the threshold point through $\mathcal{I}_{q_A}(W_t) = 1$, it follows that for $W_A > W_t$ and $W_B > W_t$

$$S_{q_A}(A \cup B) > S_{q_A}(A) + S_{q_A}(B). \quad (6.19)$$

In the same way, for $W_A < W_t$ and $W_B < W_t$, we obtain

$$S_{q_A}(A \cup B) < S_{q_A}(A) + S_{q_A}(B). \quad (6.20)$$

Temperature and pressure are given by

$$\frac{1}{T} = \frac{1}{\mathcal{I}_{q_A} - \tilde{q}_A S_{q_A}} \frac{\partial S_{q_A}}{\partial E}, \quad (6.21)$$

$$P = \frac{T}{\mathcal{I}_{q_A} - \tilde{q}_A S_{q_A}} \frac{\partial S_{q_A}}{\partial V}, \quad (6.22)$$

respectively, where $\tilde{q}_A = (q_A^{1/2} - q_A^{-1/2})^2/2$. They reduce to the standard definition of temperature and pressure in the $q_A \rightarrow 1$ limit.

The thermodynamic stability conditions now read

$$\frac{\partial^2 S_{q_A}}{\partial E^2} < \mathcal{A}_{q_A} \left(\frac{\partial S_{q_A}}{\partial E} \right)^2, \quad (6.23)$$

$$\frac{\partial^2 S_{q_A}}{\partial E^2} \frac{\partial^2 S_{q_A}}{\partial V^2} - \left(\frac{\partial^2 S_{q_A}}{\partial E \partial V} \right)^2 > \mathcal{A}_{q_A} \mathcal{B}_{q_A}, \quad (6.24)$$

with

$$\mathcal{A}_{q_A} = 2 \tilde{q}_A \frac{(\tilde{q}_A + 1) S_{q_A} - \mathcal{I}_{q_A}}{(\mathcal{I}_{q_A} - \tilde{q}_A S_{q_A})^2}. \quad (6.25)$$

The sign of Eq. (6.25) changes at the point

$$\widetilde{W}_t = q_A^{2/(q_A - q_A^{-1})}, \quad (6.26)$$

so that $\mathcal{A}_{q_A} < 0$ for $W < \widetilde{W}_t$ and $\mathcal{A}_{q_A} > 0$ for $W > \widetilde{W}_t$. On the other hand, observing that $\widetilde{W}_t < W_t$, it follows that for super-additive systems with $W_A > W_t$ and $W_B > W_t$, the concavity conditions imply the thermodynamic stability conditions.

It is worthy to observe that by posing $r = 1 - \sqrt{1 + \kappa^2} < 0$ and $q_A = \sqrt{1 + \kappa^2} - |\kappa|$ we obtain another family of entropies embodies the symmetry $q \leftrightarrow 1/q$ given by

$$S_{q_A}^*(W) = - \sum_{i=1}^W \frac{p_i^{2-q_A^{-1}} - p_i^{2-q_A}}{q_A - q_A^{-1}} = - \sum_{i=1}^W p_i \ln_{q_A}^*(p_i), \quad (6.27)$$

with $1/2 < q_A \leq 2$, where now

$$\ln_{q_A}^*(x) = \frac{x^{1-q_A^{-1}} - x^{1-q_A}}{q_A - q_A^{-1}}. \quad (6.28)$$

$\ln_{q_A}(x)$ and $\ln_{q_A}^*(x)$ are related in [16]

$$\ln_{q_A}(x) = -\ln_{q_A}^*\left(\frac{1}{x}\right), \quad (6.29)$$

and the entropies (6.15) and (6.27) are dual each other. In the microcanonical picture Eq. (6.27) becomes

$$S_{q_A}^*(W) = -\ln_{q_A}^*\left(\frac{1}{W}\right), \quad (6.30)$$

and because now the function $\mathcal{I}_{q_A}^* > 1$, the entropy (6.30) describes super-additive systems: $S_{q_A}^*(A \cup B) > S_{q_A}^*(A) + S_{q_A}^*(B)$. The function $\mathcal{A}_{q_A}^* > 0$ and the concavity conditions for the entropy (6.30) are enough to guarantee the thermodynamic stability conditions of the system for any values of the deformation parameter.

C. Kaniadakis entropy

As a last example, we discuss the entropic form introduced previously in Ref. [29] which follows from Eq. (1.6) after posed $r = 0$:

$$S_\kappa = - \sum_{i=1}^W \frac{p_i^\kappa - p_i^{-\kappa}}{2\kappa} = - \sum_{i=1}^W p_i \ln_{\{\kappa\}}(p_i), \quad (6.31)$$

where $|\kappa| < 1$, and the κ -logarithm $\ln_{\{\kappa\}}(x)$ is defined by

$$\ln_{\{\kappa\}}(x) = \frac{x^\kappa - x^{-\kappa}}{2\kappa}. \quad (6.32)$$

Its inverse function, the κ -exponential, is given by

$$\exp_{\{\kappa\}}(x) = \left(\sqrt{1 + \kappa^2 x^2} + \kappa x \right)^{1/\kappa}, \quad (6.33)$$

and satisfies the relation

$$\exp_{\{\kappa\}}(x) \exp_{\{\kappa\}}(-x) = 1, \quad (6.34)$$

which means that it increases for $x \rightarrow \infty$ and decreases for $x \rightarrow -\infty$ with the same steepness.

Remarkably, the κ -logarithm and the κ -exponential fulfill the two following mathematical proprieties

$$\exp_{\{\kappa\}}(x) \exp_{\{\kappa\}}(y) = \exp_{\{\kappa\}}(x \oplus y), \quad (6.35)$$

$$\ln_{\{\kappa\}}(xy) = \ln_{\{\kappa\}}(x) \oplus \ln_{\{\kappa\}}(y), \quad (6.36)$$

where the κ -deformed sum is defined in [50]

$$x \oplus y = x \sqrt{1 + \kappa^2 y^2} + y \sqrt{1 + \kappa^2 x^2}. \quad (6.37)$$

Eqs. (6.35) and (6.36) reduce, in the $\kappa \rightarrow 0$ limit, to the well known relations $\exp(x) \exp(y) = \exp(x + y)$ and $\ln(xy) = \ln(x) + \ln(y)$, respectively, according to $x \oplus y = x + y$.

In Ref. [30] it has been shown that the κ -sum emerges naturally within the Einstein special relativity. In fact, following the same argument presented in [30] it is possible to link the relativistic sum of the velocities

$$v_1 \oplus v_2 = \frac{v_1 + v_2}{1 + v_1 v_2 / c^2}, \quad (6.38)$$

with the κ -sum (6.37) in the sense of

$$p(v_1) \oplus p(v_2) = p(v_1 \oplus v_2), \quad (6.39)$$

where $\kappa = 1/mc$ and $p(v) = mv/\sqrt{1 - v^2/c^2}$ is the relativistic momentum of a particle with rest mass m . In this way, the origin of the κ -deformation are related to the finite value of light speed c . In particular, in the classical limit $c \rightarrow \infty$ the parameter κ approaches to zero and the κ -entropy reduces to the Boltzmann-Gibbs one.

In agreement with this interpretation, we consider statistical systems (physical or not) that can achieve an equilibrium configuration through an exchange of information between the parts of the system, propagating with a limiting velocity, like the light speed in the relativity theory. For these systems it is reasonable suppose that a mechanism similar to the one describe above, in the framework of the special relativity, can arise, so that the κ -deformation can occurs. In this way, the κ -entropy can be successfully applied in the study of the statistical proprieties of these systems.

An important physical example where the κ -distribution has been successfully applied is in the reproduction of the energy distribution of the fluxes of cosmic rays [30] (see also [51]). Moreover the κ -entropy has been applied in the study of the fracture propagation in brittle material, showing a good accordance with the results obtained experimentally and with the ones obtained through numerical simulations [52].

Finally, we recall that, like the previous one-parameter

deformed entropic forms, also the κ -entropy fulfils many physically relevant proprieties. In particular, in Ref. [53], it has been shown its experimental stability whereas in Ref. [49] the finite entropy production in time unit in connection with the Pesin identity for a system described by the entropy (6.31) has been discussed.

In the microcanonical picture, Eq. (6.31) becomes

$$S_\kappa = \ln_{\{\kappa\}}(W), \quad (6.40)$$

and we introduce the function $\mathcal{I}_\kappa$ that, according to Eq. (3.7), can be written in

$$\mathcal{I}_\kappa = \sqrt{1 + \kappa^2 S_\kappa^2}, \quad (6.41)$$

so that $\mathcal{I}_\kappa \geq 1$. As a consequence Eq. (3.3) becomes

$$S_\kappa(A \cup B) = S_\kappa(A) \sqrt{1 + \kappa^2 [S_\kappa(B)]^2} + S_\kappa(B) \sqrt{1 + \kappa^2 [S_\kappa(A)]^2}, \quad (6.42)$$

which can be also written in

$$S_\kappa(A \cup B) = S_\kappa(A) \oplus S_\kappa(B), \quad (6.43)$$

according to Eq. (6.37). Eqs. (6.42) or (6.43) imply the relation

$$S_\kappa(A \cup B) > S_\kappa(A) + S_\kappa(B), \quad (6.44)$$

stating that the κ -entropy in the microcanonical picture is always super-additive.

Temperature and pressure are given by

$$\frac{1}{T} = \frac{1}{\sqrt{1 + \kappa^2 S_\kappa^2}} \frac{\partial S_\kappa}{\partial E}, \quad (6.45)$$

$$P = \frac{T}{\sqrt{1 + \kappa^2 S_\kappa^2}} \frac{\partial S_\kappa}{\partial V}, \quad (6.46)$$

respectively, and reduce to the standard definitions in the $\kappa \rightarrow 0$ limit.

Finally, the thermodynamic stability conditions become [19, 20]

$$\frac{\partial^2 S_\kappa}{\partial E^2} < \mathcal{A}_\kappa \left(\frac{\partial S_\kappa}{\partial E} \right)^2, \quad (6.47)$$

$$\frac{\partial^2 S_\kappa}{\partial E^2} \frac{\partial^2 S_\kappa}{\partial V^2} - \left(\frac{\partial^2 S_\kappa}{\partial E \partial V} \right)^2 > \mathcal{A}_\kappa \mathcal{B}_\kappa, \quad (6.48)$$

where

$$\mathcal{A}_\kappa = \frac{\kappa^2 S_\kappa}{1 + \kappa^2 S_\kappa^2}. \quad (6.49)$$

The function (6.49) is always positive and, as a consequence, the concavity conditions for the entropy (6.40) are enough to guarantee the thermodynamic stability conditions of the system for any values of the deformation parameter.

VII. CONCLUDING REMARKS

In the present work we have investigated the thermodynamic stability conditions for a microcanonical system described by the Sharma-Taneja-Mittal entropy, and their relation with the concavity conditions for this entropy.

The main results can be summarized in the following two points:

Firstly, we have analyzed the “composability” rule for statistically independent systems described by the entropy (2.6). It has been shown that the parameter space $\mathcal{R}$ can be split into two disjoint regions. In the region $\mathcal{R}|_{r \leq 0}$ the entropy $S_{\kappa, r}$ shows a super-additivity behavior: $S_{\kappa, r}(A \cup B) > S_{\kappa, r}(A) + S_{\kappa, r}(B)$. Otherwise, in the region $\mathcal{R}|_{r > 0}$, the behavior of the entropy is not well defined, depending on the *size* of the two systems A and B. In particular it has been shown that, given the threshold point $W_t(\kappa, r) > 1$, when the *size* of the two parts A and B are smaller than W_t , in the sense of $W_A < W_t$ and $W_B < W_t$, the system exhibits a sub-additive behavior $S_{\kappa, r}(A \cup B) < S_{\kappa, r}(A) + S_{\kappa, r}(B)$, becoming super-additive when both $W_A > W_t$ and $W_B > W_t$.

Secondly, we have inquired on the thermodynamic stability conditions of the equilibrium configuration. In the Boltzmann theory the concavity conditions imply the thermodynamic stability conditions. Such a situation changes when the system is described by the entropy $S_{\kappa, r}$. We have shown that, starting from an equilibrium configuration of the system $A \cup B$, and supposed an exchange of a small quantity of heat and/or work between the two parts A and B, assumed statistically independent, if the entropy of the system $S_{\kappa, r}(A \cup B)$ is larger than the sum of the entropy of the two systems $S_{\kappa, r}(A)$ and $S_{\kappa, r}(B)$, the concavity conditions still imply the thermodynamic stability conditions. In the opposite situation, in spite of the concavity of $S_{\kappa, r}$, stability requires *large* systems, in the sense of $W_A > W_t$ and $W_B > W_t$.

APPENDIX A.

In this Appendix we summarize some mathematical properties of the deformed logarithm [31, 32]

$$\ln_{\{\kappa, r\}}(x) = \frac{x^{r+\kappa} - x^{r-\kappa}}{2\kappa}. \quad (\text{A. 1})$$

Let $\mathcal{R}$ be the region in the parametric space, defined by

$$\mathbb{R}^2 \supset \mathcal{R} = \begin{cases} -|\kappa| \leq r \leq |\kappa| & \text{if } 0 \leq |\kappa| < \frac{1}{2}, \\ |\kappa| - 1 < r < 1 - |\kappa| & \text{if } \frac{1}{2} \leq |\kappa| < 1. \end{cases} \quad (\text{A. 2})$$

For any $(\kappa, r) \in \mathcal{R}$, the $\ln_{\{\kappa, r\}}(x) = \ln_{\{-\kappa, r\}}(x)$, is a continuous, monotonic, increasing and concave function for $x \in \mathbb{R}^+$, with $\ln_{\{\kappa, r\}}(\mathbb{R}^+) \subseteq \mathbb{R}$, fulfilling the relation $\int_0^1 \ln_{\{\kappa, r\}}(x^{\pm 1}) dx = \mp 1 / [(1 \pm r)^2 - \kappa^2]$. The standard logarithm is recovered in the $(\kappa, r) \rightarrow (0, 0)$ limit:

$$\ln_{\{0, 0\}}(x) = \ln(x).$$

Eq. (A. 1) satisfies the relation

$$\ln_{\{\kappa, r\}}(x) = -\ln_{\{\kappa, -r\}}\left(\frac{1}{x}\right), \quad (\text{A. 3})$$

and for $r = 0$, reproduces the well known properties of the standard logarithm $\ln_{\{\kappa, 0\}}(x) = -\ln_{\{\kappa, 0\}}(1/x)$. From Eq. (A. 3) it follows that the behavior of $\ln_{\{\kappa, r\}}(x)$ for $r > 0$ and $0 \leq x \leq 1$ is related to the one for $r < 0$ and $x > 1$.

We observe that the deformed logarithm is a solution of the following differential-functional equation

$$\frac{d}{dx} [x \ln_{\{\kappa, r\}}(x)] = \lambda \ln_{\{\kappa, r\}}\left(\frac{x}{\alpha}\right), \quad (\text{A. 4})$$

with the following boundary conditions $\ln_{\{\kappa, r\}}(1) = 0$ and $d \ln_{\{\kappa, r\}}(x)/dx|_{x=1} = 1$ and the constants α and λ given in Eqs. (1.4)-(1.5).

The inverse function of Eq. (A. 1), namely the deformed exponential $\exp_{\{\kappa, r\}}(x)$, exists for any $(\kappa, r) \in \mathcal{R}$, since $\ln_{\{\kappa, r\}}(x)$ is a strictly monotonic function. Its analytical properties are well-defined and follow from the corresponding ones of the deformed logarithm. Nevertheless, the explicit expression of the $\exp_{\{\kappa, r\}}(x)$ can be obtained only for particular relationships between r and κ .

By inspection there exists a point $x_0(\kappa, r)$ such that the inequality $\ln_{\{\kappa, r\}}(x) \geq \ln x$ holds for $x > x_0(\kappa, r)$. We see that $x_0(\kappa, r)$ is a monotonic decreasing function w.r.t r with $x_0(\kappa, r) > 1$ for $r < 0$ and $x_0(\kappa, r) < 1$ for $r > 0$. In particular $x_0(\kappa, -\kappa) = +\infty$, $x_0(\kappa, 0) = 1$ and $x_0(\kappa, \kappa) = 0$.

Entropy $S_{\kappa, r}(p)$, introduced in Eq. (1.6), is related to the function $\ln_{\{\kappa, r\}}(x)$ through the relation

$$S_{\kappa, r}(p) = -\sum_{i=1}^W p_i \ln_{\{\kappa, r\}}(p_i). \quad (\text{A. 5})$$

The deformed logarithm (A. 1) can be obtained from the two-parametric generalization of the Jackson derivative, previously proposed in Ref. [54]

$$\frac{d_{\kappa, r} f(x)}{d_{\kappa, r} x} = \frac{f((r+\kappa)x) - f((r-\kappa)x)}{2\kappa x}, \quad (\text{A. 6})$$

by posing

$$\ln_{\{\kappa, r\}}(x) = \frac{d_{\kappa, r} x^y}{d_{\kappa, r} y} \Big|_{y=1}. \quad (\text{A. 7})$$

Some properties of the deformed logarithm can be naturally understood as those of the generalized Jackson derivative (A. 6). For instance, from the generalized Leibnitz rule

$$\begin{aligned} \frac{d_{\kappa, r} f(x) g(x)}{d_{\kappa, r} x} &= \frac{d_{\kappa, r} f(x)}{d_{\kappa, r} x} g((r+\kappa)x) \\ &+ f((r-\kappa)x) \frac{d_{\kappa, r} g(x)}{d_{\kappa, r} x}, \end{aligned} \quad (\text{A. 8})$$

we obtain the following useful relation

$$\ln_{\{\kappa, r\}}(xy) = x^{r+\kappa} \ln_{\{\kappa, r\}}(y) + y^{r-\kappa} \ln_{\{\kappa, r\}}(x), \quad (\text{A. 9})$$

and by using the identity $y^{r-\kappa} = y^{r+\kappa} - 2\kappa \ln_{\{\kappa, r\}}(y)$, Eq. (A. 9) becomes

$$\begin{aligned} \ln_{\{\kappa, r\}}(xy) &= x^{r+\kappa} \ln_{\{\kappa, r\}}(y) + y^{r+\kappa} \ln_{\{\kappa, r\}}(x) \\ &\quad - 2\kappa \ln_{\{\kappa, r\}}(x) \ln_{\{\kappa, r\}}(y). \end{aligned} \quad (\text{A. 10})$$

Moreover, recalling the $\kappa \leftrightarrow -\kappa$ symmetry, Eq. (A. 10) can be rewritten in

$$\begin{aligned} \ln_{\{\kappa, r\}}(xy) &= u_{\{\kappa, r\}}(x) \ln_{\{\kappa, r\}}(y) \\ &\quad + u_{\{\kappa, r\}}(x) \ln_{\{\kappa, r\}}(y), \end{aligned} \quad (\text{A. 11})$$

where we have introduced the function

$$u_{\{\kappa, r\}}(x) = \frac{x^{r+\kappa} + x^{r-\kappa}}{2}. \quad (\text{A. 12})$$

For any $(\kappa, r) \in \mathcal{R}$ the function $u_{\{\kappa, r\}}(x) = u_{\{-\kappa, r\}}(x)$, is continuous for $x \in \mathbb{R}^+$, with $u_{\{\kappa, r\}}(\mathbb{R}^+) \subseteq \mathbb{R}^+$, $u_{\{\kappa, r\}}(0) = u_{\{\kappa, r\}}(+\infty) = +\infty$ for $r \neq |\kappa|$, satisfies the relation $u_{\{\kappa, r\}}(x) = u_{\{\kappa, -r\}}(1/x)$ and reduces to the unity in the $(\kappa, r) \rightarrow (0, 0)$ limit: $u_{\{0, 0\}}(x) = 1$. It reaches the minimum values

$$u_{\{\kappa, r\}}(x_m) = \kappa \frac{(\kappa - r)^{(r-\kappa)/2\kappa}}{(\kappa + r)^{(r+\kappa)/2\kappa}}, \quad (\text{A. 13})$$

at

$$x_m = \left(\frac{\kappa - r}{\kappa + r} \right)^{1/2\kappa}. \quad (\text{A. 14})$$

In particular, for any $(\kappa, r) \in \mathcal{R}|_{r \leq 0}$, $x_m \geq 1$, and taking into account that $u_{\{\kappa, r\}}(1) = 1$, it follows $1 \leq u_{\{\kappa, r\}}(x) \leq \infty$ when $x \in (0, 1)$. For any $(\kappa, r) \in \mathcal{R}|_{r > 0}$, from Eq. (A. 14) we obtain $0 \leq x_m \leq 1$ and from Eq. (A. 13) it follows $1/2 \leq u_{\{\kappa, r\}}(x_m) \leq 1$.

By inspection, it follows that there exist a threshold point $x_t(\kappa, r)$, defined by $u_{\{\kappa, r\}}(x_t) = 1$. $x_t(\kappa, r)$ is monotonic decreasing function w.r.t. r , with $x_t(\kappa, -\kappa) = +\infty$, $x_t(\kappa, 0) = 1$ and $x_t(\kappa, \kappa) = 0$, such that $1/2 \leq u_{\{\kappa, r\}}(x_m) \leq u_{\{\kappa, r\}}(x) \leq 1$ for $x \geq x_t(\kappa, r)$ and $1 \leq u_{\{\kappa, r\}}(x) \leq +\infty$ for $0 \leq x \leq x_t(\kappa, r)$.

Finally, we remark that the function (A. 12) fulfills the following relation

$$u_{\{\kappa, r\}}(xy) = u_{\{\kappa, r\}}(x) u_{\{\kappa, r\}}(y) + \kappa^2 \ln_{\{\kappa, r\}}(x) \ln_{\{\kappa, r\}}(y). \quad (\text{A. 15})$$

We observe that, like the deformed logarithm, the function $u_{\{\kappa, r\}}(x)$ is a solution of the differential-functional equation (1.2) with the following boundary conditions $u_{\{\kappa, r\}}(1) = 1$ and $du_{\{\kappa, r\}}(x)/dx|_{x=1} = r$ and the constants α and λ given in Eqs. (1.4)-(1.5). Moreover the

two functions $\ln_{\{\kappa, r\}}(x)$ and $u_{\{\kappa, r\}}(x)$ are related by the relation

$$u_{\{\kappa, r\}}(x) = x^{r+\kappa} - \kappa \ln_{\{\kappa, r\}}(x). \quad (\text{A. 16})$$

The function $\mathcal{I}_{\kappa, r}(p)$, introduced in Eq. (3.6), is related to the function $u_{\{\kappa, r\}}(x)$ through the relation

$$\mathcal{I}_{\kappa, r}(p) = \sum_{i=1}^W p_i u_{\{\kappa, r\}}(p_i). \quad (\text{A. 17})$$

From the definitions (A. 5), (A. 17) and Eqs. (A. 11), (A. 15), we obtain the useful relations

$$S_{\kappa, r}(A \cup B) = S_{\kappa, r}(A) \mathcal{I}_{\kappa, r}(B) + \mathcal{I}_{\kappa, r}(A) S_{\kappa, r}(B), \quad (\text{A. 18})$$

$$\mathcal{I}_{\kappa, r}(A \cup B) = \mathcal{I}_{\kappa, r}(A) \mathcal{I}_{\kappa, r}(B) + \kappa^2 S_{\kappa, r}(A) S_{\kappa, r}(B), \quad (\text{A. 19})$$

stating the additivity of $S_{\kappa, r}$ and $\mathcal{I}_{\kappa, r}$ for statistical independent systems $p^{A \cup B} = \{p_i^A p_j^B\}$ with $i = 1, \dots, W_A$ and $j = 1, \dots, W_B$.

APPENDIX B.

In this Appendix we derive the equilibrium conditions given in Eqs. (4.4)-(4.5), and the thermodynamic stability conditions given in Eqs. (5.7)-(5.8).

Let us suppose that the whole system $A \cup B$ initially at equilibrium undergoes a small transfer of heat and/or work between the two parts A and B, with the constraints

$$\delta(E_A + E_B) = 0, \quad (\text{B. 1})$$

$$\delta(V_A + V_B) = 0. \quad (\text{B. 2})$$

Recalling that the entropy evaluated at an equilibrium configuration is a maximum, we have

$$S_{\kappa, r}(A \cup B) > S_{\kappa, r}((A + \delta A) \cup (B + \delta B)), \quad (\text{B. 3})$$

and taking into account of Eq. (A. 18) it follows

$$\begin{aligned} &S_{\kappa, r}(A) \mathcal{I}_{\kappa, r}(B) + \mathcal{I}_{\kappa, r}(A) S_{\kappa, r}(B) > \\ &S_{\kappa, r}(A + \delta A) \mathcal{I}_{\kappa, r}(B + \delta B) \\ &+ \mathcal{I}_{\kappa, r}(A + \delta A) S_{\kappa, r}(B + \delta B). \end{aligned} \quad (\text{B. 4})$$

We expand the r.h.s of Eq. (B. 4) up to the second order in δE and δV , where $\delta E \equiv \delta E_A = -\delta E_B$ and $\delta V \equiv \delta V_A = -\delta V_B$. According to the MaxEnt principle, the first order terms must vanish

$$\begin{aligned} &\left(\frac{\partial S_{\kappa, r}(A)}{\partial E_A} \mathcal{I}_{\kappa, r}(B) + \frac{\partial \mathcal{I}_{\kappa, r}(A)}{\partial E_A} S_{\kappa, r}(B) \right. \\ &\quad \left. - S_{\kappa, r}(A) \frac{\partial \mathcal{I}_{\kappa, r}(B)}{\partial E_B} - \mathcal{I}_{\kappa, r}(A) \frac{\partial S_{\kappa, r}(B)}{\partial E_B} \right) \delta E \\ &+ \left(\frac{\partial S_{\kappa, r}(A)}{\partial V_A} \mathcal{I}_{\kappa, r}(B) + \frac{\partial \mathcal{I}_{\kappa, r}(A)}{\partial V_A} S_{\kappa, r}(B) \right. \\ &\quad \left. - S_{\kappa, r}(A) \frac{\partial \mathcal{I}_{\kappa, r}(B)}{\partial V_B} - \mathcal{I}_{\kappa, r}(A) \frac{\partial S_{\kappa, r}(B)}{\partial V_B} \right) \delta V = 0. \end{aligned} \quad (\text{B. 5})$$

By using the relations

$$\frac{\partial \mathcal{I}_{\kappa, r}}{\partial X} = \frac{\kappa^2 S_{\kappa, r} - r \mathcal{I}_{\kappa, r}}{\mathcal{I}_{\kappa, r} - r S_{\kappa, r}} \frac{\partial S_{\kappa, r}}{\partial X}, \quad (\text{B. 6})$$

with $X \equiv E$ or $X \equiv V$, Eq. (B. 5) becomes

$$\begin{aligned} & \left(\mathcal{I}_{\kappa, r} - r S_{\kappa, r} \right) \Big|_{\text{AUB}} \left[\frac{1}{\mathcal{I}_{\kappa, r} - r S_{\kappa, r}} \frac{\partial S_{\kappa, r}}{\partial E} \Big|_{\text{A}} \delta E \right. \\ & + \frac{1}{\mathcal{I}_{\kappa, r} - r S_{\kappa, r}} \frac{\partial S_{\kappa, r}}{\partial V} \Big|_{\text{A}} \delta V - \frac{1}{\mathcal{I}_{\kappa, r} - r S_{\kappa, r}} \frac{\partial S_{\kappa, r}}{\partial E} \Big|_{\text{B}} \delta E \\ & \left. - \frac{1}{\mathcal{I}_{\kappa, r} - r S_{\kappa, r}} \frac{\partial S_{\kappa, r}}{\partial V} \Big|_{\text{B}} \delta V \right] = 0, \end{aligned} \quad (\text{B. 7})$$

where we use the relations (A. 18) and (A. 19). Taking into account that

$$\mathcal{I}_{\kappa, r} \left(\frac{1}{W} \right) - r S_{\kappa, r} \left(\frac{1}{W} \right) \quad (\text{B. 8})$$

$$= \frac{1}{W} \frac{d}{d(1/W)} \ln_{\{\kappa, r\}} \left(\frac{1}{W} \right) > 0, \quad (\text{B. 9})$$

through Eq. (B. 7) the two equilibrium conditions follow

$$\frac{1}{\mathcal{I}_{\kappa, r} - r S_{\kappa, r}} \frac{\partial S_{\kappa, r}}{\partial E} \Big|_{\text{A}} = \frac{1}{\mathcal{I}_{\kappa, r} - r S_{\kappa, r}} \frac{\partial S_{\kappa, r}}{\partial E} \Big|_{\text{B}}, \quad (\text{B. 10})$$

$$\frac{1}{\mathcal{I}_{\kappa, r} - r S_{\kappa, r}} \frac{\partial S_{\kappa, r}}{\partial V} \Big|_{\text{A}} = \frac{1}{\mathcal{I}_{\kappa, r} - r S_{\kappa, r}} \frac{\partial S_{\kappa, r}}{\partial V} \Big|_{\text{B}}, \quad (\text{B. 11})$$

which coincide with Eqs. (4.4) and (4.5).

In order to obtain the thermodynamic stability conditions let us consider the second order terms in the expansion of Eq. (B. 4)

$$\begin{aligned} & \frac{1}{2} \left[\mathcal{I}_{\kappa, r}(\text{A}) \frac{\partial^2 S_{\kappa, r}(\text{B})}{\partial E_{\text{B}}^2} - 2 \frac{\partial \mathcal{I}_{\kappa, r}(\text{A})}{\partial E_{\text{A}}} \frac{S_{\kappa, r}(\text{B})}{\partial E_{\text{B}}} \right. \\ & + \frac{\partial^2 \mathcal{I}_{\kappa, r}(\text{A})}{\partial E_{\text{A}}^2} S_{\kappa, r}(\text{B}) + \mathcal{I}_{\kappa, r}(\text{B}) \frac{\partial^2 S_{\kappa, r}(\text{A})}{\partial E_{\text{A}}^2} \\ & \left. - 2 \frac{\partial \mathcal{I}_{\kappa, r}(\text{B})}{\partial E_{\text{B}}} \frac{S_{\kappa, r}(\text{A})}{\partial E_{\text{A}}} + \frac{\partial^2 \mathcal{I}_{\kappa, r}(\text{B})}{\partial E_{\text{B}}^2} S_{\kappa, r}(\text{A}) \right] (\delta E)^2 \\ & + \frac{1}{2} \left[\mathcal{I}_{\kappa, r}(\text{A}) \frac{\partial^2 S_{\kappa, r}(\text{B})}{\partial V_{\text{B}}^2} - 2 \frac{\partial \mathcal{I}_{\kappa, r}(\text{A})}{\partial V_{\text{A}}} \frac{S_{\kappa, r}(\text{B})}{\partial V_{\text{B}}} \right. \\ & + \frac{\partial^2 \mathcal{I}_{\kappa, r}(\text{A})}{\partial V_{\text{A}}^2} S_{\kappa, r}(\text{B}) + \mathcal{I}_{\kappa, r}(\text{B}) \frac{\partial^2 S_{\kappa, r}(\text{A})}{\partial V_{\text{A}}^2} \\ & \left. - 2 \frac{\partial \mathcal{I}_{\kappa, r}(\text{B})}{\partial V_{\text{B}}} \frac{S_{\kappa, r}(\text{A})}{\partial V_{\text{A}}} + \frac{\partial^2 \mathcal{I}_{\kappa, r}(\text{B})}{\partial V_{\text{B}}^2} S_{\kappa, r}(\text{A}) \right] (\delta V)^2 \\ & + \left[\mathcal{I}_{\kappa, r}(\text{A}) \frac{\partial^2 S_{\kappa, r}(\text{B})}{\partial E_{\text{B}} \partial V_{\text{B}}} - \frac{\partial \mathcal{I}_{\kappa, r}(\text{A})}{\partial V_{\text{A}}} \frac{\partial S_{\kappa, r}(\text{B})}{\partial E_{\text{B}}} \right. \end{aligned}$$

$$\begin{aligned} & - \frac{\partial \mathcal{I}_{\kappa, r}(\text{A})}{\partial E_{\text{A}}} \frac{\partial S_{\kappa, r}(\text{B})}{\partial V_{\text{B}}} + \frac{\partial^2 \mathcal{I}_{\kappa, r}(\text{A})}{\partial E_{\text{A}} \partial V_{\text{A}}} S_{\kappa, r}(\text{B}) \\ & + \mathcal{I}_{\kappa, r}(\text{B}) \frac{\partial^2 S_{\kappa, r}(\text{A})}{\partial E_{\text{A}} \partial V_{\text{A}}} - \frac{\partial \mathcal{I}_{\kappa, r}(\text{B})}{\partial V_{\text{B}}} \frac{\partial S_{\kappa, r}(\text{A})}{\partial E_{\text{A}}} \\ & \left. - \frac{\partial \mathcal{I}_{\kappa, r}(\text{B})}{\partial E_{\text{B}}} \frac{\partial S_{\kappa, r}(\text{A})}{\partial V_{\text{A}}} + \frac{\partial^2 \mathcal{I}_{\kappa, r}(\text{B})}{\partial E_{\text{B}} \partial V_{\text{B}}} S_{\kappa, r}(\text{A}) \right] \delta E \delta V < 0. \end{aligned} \quad (\text{B. 12})$$

By using the relation

$$\begin{aligned} \frac{\partial^2 \mathcal{I}_{\kappa, r}}{\partial X \partial Y} &= \frac{\kappa^2 S_{\kappa, r} - r \mathcal{I}_{\kappa, r}}{\mathcal{I}_{\kappa, r} - r S_{\kappa, r}} \frac{\partial^2 S_{\kappa, r}}{\partial X \partial Y} \\ &+ (\kappa^2 - r^2) \frac{\mathcal{I}_{\kappa, r}^2 - \kappa^2 S_{\kappa, r}^2}{(\mathcal{I}_{\kappa, r} - r S_{\kappa, r})^3} \frac{\partial S_{\kappa, r}}{\partial X} \frac{\partial S_{\kappa, r}}{\partial Y}, \end{aligned} \quad (\text{B. 13})$$

Eq. (B. 12) becomes

$$\begin{aligned} & \frac{1}{2} \left(\mathcal{I}_{\kappa, r} - r S_{\kappa, r} \right) \Big|_{\text{AUB}} \\ & \times \left[\frac{\mathcal{S}_{\text{EE}} (\delta E)^2 + 2 \mathcal{S}_{\text{EV}} \delta E \delta V + \mathcal{S}_{\text{VV}} (\delta V)^2}{\mathcal{I}_{\kappa, r} - r S_{\kappa, r}} \Big|_{\text{A}} \right. \\ & \left. + \frac{\mathcal{S}_{\text{EE}} (\delta E)^2 + 2 \mathcal{S}_{\text{EV}} \delta E \delta V + \mathcal{S}_{\text{VV}} (\delta V)^2}{\mathcal{I}_{\kappa, r} - r S_{\kappa, r}} \Big|_{\text{B}} \right] < 0, \end{aligned} \quad (\text{B. 14})$$

where we have defined

$$\mathcal{S}_{\text{XY}} = \frac{\partial^2 S_{\kappa, r}}{\partial X \partial Y} - \frac{(\kappa^2 + r^2) S_{\kappa, r} - 2 r \mathcal{I}_{\kappa, r}}{(\mathcal{I}_{\kappa, r} - r S_{\kappa, r})^2} \frac{\partial S_{\kappa, r}}{\partial X} \frac{\partial S_{\kappa, r}}{\partial Y}, \quad (\text{B. 15})$$

and taking into account Eq. (B. 9), from Eq. (B. 14) it follows that the inequality

$$\mathcal{S}_{\text{EE}} (\delta E)^2 + 2 \mathcal{S}_{\text{EV}} \delta E \delta V + \mathcal{S}_{\text{VV}} (\delta V)^2 < 0, \quad (\text{B. 16})$$

must holds for both systems A and B.

In Eq. (B. 16), (by posing $\delta V = 0$ and $\delta E = 0$, respectively), it follows

$$\mathcal{S}_{\text{EE}} < 0, \quad \mathcal{S}_{\text{VV}} < 0, \quad (\text{B. 17})$$

and, multiplying Eq. (B. 16) by $\mathcal{S}_{\text{EE}}$, we obtain

$$(\mathcal{S}_{\text{EE}} \delta E + \mathcal{S}_{\text{EV}} \delta V)^2 + (\mathcal{S}_{\text{EE}} \mathcal{S}_{\text{VV}} - \mathcal{S}_{\text{EV}}^2) (\delta V)^2 > 0, \quad (\text{B. 18})$$

which implies

$$\mathcal{S}_{\text{EE}} \mathcal{S}_{\text{VV}} - \mathcal{S}_{\text{EV}}^2 > 0. \quad (\text{B. 19})$$

Eqs. (B. 17) and (B. 19) are the thermodynamic stability conditions.

In particular, Eqs. (B. 17) can be written in

$$\frac{\partial^2 S_{\kappa, r}}{\partial X^2} < \mathcal{A}_{\kappa, r} \left(\frac{\partial S_{\kappa, r}}{\partial X} \right)^2, \quad (\text{B. 20})$$

with

$$\mathcal{A}_{\kappa, r} = \frac{(\kappa^2 + r^2) S_{\kappa, r} - 2 r \mathcal{I}_{\kappa, r}}{(\mathcal{I}_{\kappa, r} - r S_{\kappa, r})^2}, \quad (\text{B. 21})$$

whilst from Eq. (B. 19) we obtain

$$\frac{\partial^2 S_{\kappa, r}}{\partial E^2} \frac{\partial^2 S_{\kappa, r}}{\partial V^2} - \left(\frac{\partial^2 S_{\kappa, r}}{\partial E \partial V} \right)^2 > \mathcal{A}_{\kappa, r} \mathcal{B}_{\kappa, r}, \quad (\text{B. 22})$$

where

$$\mathcal{B}_{\kappa, r} = \left(\frac{\partial^2 S_{\kappa, r}}{\partial E^2} \right)^{-1} \left\{ \left(\frac{\partial^2 S_{\kappa, r}}{\partial E^2} \frac{\partial S_{\kappa, r}}{\partial V} - \frac{\partial^2 S_{\kappa, r}}{\partial E \partial V} \frac{\partial S_{\kappa, r}}{\partial E} \right)^2 + \left(\frac{\partial S_{\kappa, r}}{\partial E} \right)^2 \left[\frac{\partial^2 S_{\kappa, r}}{\partial E^2} \frac{\partial^2 S_{\kappa, r}}{\partial V^2} - \left(\frac{\partial^2 S_{\kappa, r}}{\partial E \partial V} \right)^2 \right] \right\},$$

and, according to Eq. (2.8)-(2.9), it follows $\mathcal{B}_{\kappa, r} < 0$.

-
- [1] E.T. Jaynes, “Papers on Probability, Statistics and Statistical Physics”, R.D. Rosenkrantz editors, (Kluwer Academic Publishers, Dordrecht, 1989).
 - [2] H.B. Callen, *Thermodynamics and an Introduction to Thermostatistics*, (Wiley, New York, 1985).
 - [3] D. Chandler, *Introduction to Modern Statistical Mechanics*, (Oxford University Press, Oxford, 1987).
 - [4] D.H. Zanette, and P.A. Alemany, Phys. Rev. Lett. **75**, 366 (1995).
 - [5] A. Compte, and D. Jou, J. Phys. A **29**, 4321 (1996).
 - [6] C. Beck, Physica A **277**, 115 (2000).
 - [7] J. Makino, Astrophys. J. **141**, 796 (1996).
 - [8] H. Uys, H.G. Miller, and F.C. Khanna, Phys. Lett. A **289**, 264 (2001).
 - [9] B. Tanatar, Phys. Rev. E **65**, 046105 (2002).
 - [10] A. Rossani, A.M. Scarfone, Physica A **282**, 212 (2000).
 - [11] A. Upadhyaya, J.-P. Rieu, J.A. Glazier, Y. Sawada, Physica A **293**, 549 (2001).
 - [12] S. Abe, “Nonextensive Statistical Mechanics and its Applications”, Y. Okamoto editors, (Springer 2001).
 - [13] Special issue of Physica A **305**, Nos. 1/2 (2002), edited by G. Kaniadakis, M. Lissia, and A. Rapisarda.
 - [14] Special issue of Physica A **340**, Nos. 1/3 (2004), edited by G. Kaniadakis, and M. Lissia.
 - [15] J. Naudts, Physica A **340**, 32 (2004).
 - [16] J. Naudts, Physica A **316**, 323 (2002).
 - [17] J.D. Ramshaw, Phys. Lett. A **198**, 119 (1995).
 - [18] T. Wada, Phys. Lett. A **297**, 334 (2002).
 - [19] T. Wada, Continuum Mech. Thermodyn. **16**, 263 (2004).
 - [20] T. Wada, Physica A **340**, 126 (2004).
 - [21] B.D. Sharma, and I.J. Taneja, Metrika **22**, 205 (1975).
 - [22] B.D. Sharma, and D.P. Mittal, J. Math. Sci. **10**, 28 (1975).
 - [23] D.P. Mittal, Metrika **22**, 35 (1975).
 - [24] E.P. Borges, and I. Roditi, Phys. Lett. A **246**, 399 (1998).
 - [25] C. Tsallis, J. Stat. Phys. **52**, 479 (1988).
 - [26] For a full bibliography see <http://tsallis.cat.cbpf.br/biblio.htm>
 - [27] S. Abe, Phys. Lett. A **224**, 326 (1997).
 - [28] S. Abe, Phys. Lett. A **244**, 229 (1998).
 - [29] G. Kaniadakis, Physica A **296**, 405 (2001).
 - [30] G. Kaniadakis, Phys. Rev. E **66**, 056125 (2002).
 - [31] G. Kaniadakis, M. Lissia, and A.M. Scarfone, Phys. Rev. E (2005), **71**, 046128 (2005).
 - [32] G. Kaniadakis, M. Lissia, and A.M. Scarfone, Physica A **340**, 41 (2004).
 - [33] H. Hancock, *Theory of maxima and minima*, (Dover, New York, 1960).
 - [34] S. Abe, Phys. Rev. E **63**, 061105 (2001).
 - [35] S. Abe, Physica A **305**, 62 (2002).
 - [36] S. Abe, Physica A **300**, 417 (2001).
 - [37] S. Abe, S. Martínez, F. Pennini, and A. Plastino, Phys. Lett. A **281**, 126 (2001).
 - [38] Q.A. Wang, L. Nivanen, and A. Le Méhauté, Eur. Phys. Lett. **65**, 606 (2004).
 - [39] R. Toral, Physica A **317**, 209 (2003).
 - [40] L. Velazquez, and F. Guzman, Phys. Rev. E **65**, 046134 (2002).
 - [41] E.P. Borges, Physica A **340**, 95 (2004).
 - [42] L. Nivanen, A. Le Méhauté, and Q.A. Wang, Rep. Math. Phys. **52**, 437 (2003).
 - [43] S. Abe, Phys. Rev. E **66**, 046134 (2002).
 - [44] B. Lesche, J. Stat. Phys. **27**, 419 (1982); Phys. Rev. E **70**, 017102 (2004).
 - [45] C. Tsallis, A.R. Plastino, and W.-M. Zheng, Chaos Soliton Fractals **8**, 885 (1997).
 - [46] M. Gell-Mann, and C. Tsallis, eds. “Nonextensive Entropy - Interdisciplinary applications”, (Oxford University Press, New York, 2004).
 - [47] G.R. Guerberoff, and G.A. Raggio, Phys. Lett. A **214**, 313 (1996).
 - [48] S. Abe, G. Kaniadakis, and A.M. Scarfone, J. Phys. A: Math. Gen. **37**, 10513 (2004).
 - [49] R. Tonelli, G. Mezzorani, F. Meloni, M. Lissia, and M. Curaddu, “Entropy production and Pesin identity at the onset of chaos”, arXiv:cond -mat/0412730, (submitted).
 - [50] G. Kaniadakis, and A.M. Scarfone, Physica A **305**, 69 (2002).
 - [51] C. Tsallis, J.C. Anjos, and E.P. Borges, Phys. Lett. A **310**, 372 (2003).
 - [52] M. Cravero, G. Iabichino, G. Kaniadakis, E. Miraldi, and A.M. Scarfone, Physica A **340**, 410 (2004).
 - [53] G. Kaniadakis, and A.M. Scarfone, Physica A **340**, 102 (2004).
 - [54] R. Chakrabarti, and R. Jagannathan, J. Phys. A **24**, L711 (1991).
 - [55] Here *size* is used to indicate the value W of the accessible states of the system.